

WORLD HEALTH STATISTICS

OVERVIEW

2019

MONITORING
HEALTH FOR THE

SDGs

S U S T A I N A B L E
D E V E L O P M E N T
G O A L S



World Health
Organization

WORLD HEALTH STATISTICS

OVERVIEW

2019

MONITORING
HEALTH FOR THE
SDGs

S U S T A I N A B L E
D E V E L O P M E N T G O A L S



World Health
Organization

CONTENTS

Foreword	v
Abbreviations	vi
1. Introduction	1
2. Life-expectancy and cause of death	2
3. Health-related SDG indicators	6
3.1 Global trends	6
3.2 Differences between countries	7
3.3 Differences between females and males	9
4. Underlying data for tracking the health-related SDGs	12
5. Conclusions	14
6. Ways forward	15
References	16

FOREWORD



For 71 years, the World Health Organization has had one vision: the highest attainable standard of health for all people in all countries. Reliable data are vital for assessing whether we are making progress toward that vision, for highlighting areas of progress and for exposing where we need to change course.

Health trends are constantly evolving, and so are health systems. No health system is perfect, and all countries have people who are left behind. It is therefore important to accurately document the state of global health and how it is changing.

The *World health statistics*, published annually since 2005, is WHO's annual snapshot of the state of the world's health. Since 2016, the World Health Statistics series has focused on monitoring progress toward the Sustainable Development Goals, and the 2019 edition contains the latest available data for the health-related SDG indicators.

The SDGs address inequality wherever it exists. The 2019 report reports on trends over time and disaggregates data by WHO region, World Bank income group, and sex to identify key inequalities. The report reveals that in low-income countries, health is frequently compromised by diseases and conditions that are preventable or treatable. Many premature deaths are associated with environmental factors or the leading underlying causes of death, such as tobacco use, unhealthy diet, physical inactivity, and unhealthy consumption of alcohol.

Although the *World health statistics 2019* tells its story with numbers, the consequences are human. In low-income countries, one woman in 41 dies from maternal causes, and each maternal death greatly affects the health of surviving family members and the resilience of the community. Not only are the risks of maternal deaths elevated by poverty, but their occurrence perpetuates the cycle of poverty in poor communities from one generation to the next.

We must be relentless in our pursuit of solutions to such human tragedies, and our responses must be informed by robust, reliable data on health risks, access to services and health outcomes. At present, many countries lack the health information systems they need to accurately monitor health trends and inform decision making. That is why WHO is creating an entirely new process for strategic policy dialogue with countries, using evidence and information strategically to drive change.

We will continue to work with countries to strengthen their information systems for health and generate better data. We will also build a modern data backbone that allows sharing and analysis of data among the three levels of the organization and countries. And we will make investments to strengthen analytical capacity at all levels, to identify trends, make projections, and support acceleration of progress toward achieving health targets.

Behind every number in these pages is a person, a family, a community and a nation. My hope is that governments, health providers, academics, civil society organizations, the media and others use these numbers to promote health, keep the world safe and serve the vulnerable.

A handwritten signature in black ink, which appears to read 'Tedros Adhanom Ghebreyesus'. The signature is fluid and cursive.

Dr Tedros Adhanom Ghebreyesus

Director-General
World Health Organization

ABBREVIATIONS

AFR	WHO African Region
AIDS	acquired immunodeficiency syndrome
AMR	WHO Region of the Americas
COPD	chronic obstructive pulmonary disease
CVD	cardiovascular disease
DTP3	diphtheria-tetanus-pertussis third dose
EMR	WHO Eastern Mediterranean Region
EUR	WHO European Region
HALE	healthy life expectancy
HI	high income
HIV	human immunodeficiency virus
LI	low income
LMI	lower-middle income
M/F	male to female (ratio)
MCV2	measles-containing vaccine second dose
NCD	noncommunicable disease
NTD	neglected tropical disease
ODA	official development assistance
PCV3	pneumococcal conjugate vaccine third dose
PPP	purchasing power parity
SDG	Sustainable Development Goal
SEAR	WHO South-East Asia Region
TB	tuberculosis
UHC	universal health coverage
UMI	upper-middle income
WB	World Bank
WHO	World Health Organization
WPR	WHO Western Pacific Region

INTRODUCTION

The World Health Statistics series is the World Health Organization's (WHO's) annual compilation of health statistics for its 194 Member States. The series is produced by WHO's Division for Data, Analytics and Delivery, in collaboration with WHO technical departments and regional offices.

World health statistics 2019 summarizes recent trends and levels in life expectancy and causes of death, and reports on the health and health-related Sustainable Development Goals (SDGs) and associated targets. Where possible, the 2019 report disaggregates data by WHO region, World Bank income group, and sex; it also discusses differences in health status and access to preventive and curative services, particularly in relation to differences between men and women. Where possible, it indicates the roles of sex as a

biological determinant, and of gender as a social construct, in accounting for the observed differences (Table O1).

The analyses presented are not exhaustive; nevertheless, it is hoped that the report will raise awareness of some critical sex and gender differences in health outcomes, highlight the importance of those differences in the attainment of the SDGs, and encourage the roles of sex and gender to be systematically taken into account when collecting data, analysing health situations, formulating policies and designing health programmes.

This overview summarizes the principal findings of the *World health statistics 2019*. The full report can be accessed at https://www.who.int/gho/publications/world_health_statistics/2019/en/.

Table O.1
Definitions of sex and gender

Sex	Gender
Sex refers to the biological characteristics that define humans as female or male. These sets of biological characteristics are not mutually exclusive, because there are individuals who are born with physical or biological sex characteristics who do not fit the traditional definitions of female or male (intersex).	Gender refers to the socially constructed norms, roles and relations of and among women, men, boys and girls. Gender also refers to expressions and identities of women, men, boys, girls and gender-diverse people. Gender is inextricable from other social and structural determinants shaping health and equity and can vary across time and place.
Sex differences influence chromosomes, gene expression, hormones, immune system and anatomy (e.g. body size, and sexual and reproductive anatomy).	Gender differences influence exposure to risk factors, health-seeking and risk-taking behaviours, access to and use of health information, and promotive, preventive, curative, rehabilitative and palliative health services.
<p>Examples of sex-specific conditions:</p> <ul style="list-style-type: none"> • cervical cancer (women); • prostate cancer (men); and • X-linked immune regulators may enhance immune responses in female children, resulting in reduced mortality among girls aged under 5 years. 	<p>Examples of gender-related factors resulting in differential health outcomes:</p> <ul style="list-style-type: none"> • child marriage increases the risks related to early pregnancy among girls; • higher rates of male employment in the transport industry exposes men to higher risks of death on the roads; • gender norms in some settings promote smoking and alcohol use among men, although young women are increasingly targeted with messages from the tobacco industry; • women's use of health services may be limited by lack of access to and control of household financial resources; • men's use of health services may be influenced by their employment conditions: also, men's limited use of health services may be influenced by socially constructed roles in which seeking health care is not seen as manly; and • transgender people often experience high rates of stigma and discrimination in health care settings, and lack appropriate services tailored to their needs.

LIFE EXPECTANCY AND CAUSE OF DEATH


Females live longer than males

In 2019, more than 141 million children will be born: 73 million boys and 68 million girls. Based on recent mortality risks the boys will live, on average, 69.8 years and the girls 74.2 years – a difference of 4.4 years. Life expectancy at age 60 years is also greater for women than men: 21.9 versus 19.0 years.

Between 2000 and 2016, global life expectancy at birth, for both sexes combined, increased by 5.5 years, from 66.5 to

72.0 years. The number of years lived in full health – that is, healthy life expectancy (HALE) – also increased over that period, from 58.5 years in 2000 to 63.3 years in 2016 (Fig. O.1). HALE is greater in women than men at birth (64.8 versus 62.0 years) and at age 60 years (16.8 versus 14.8 years). However, the number of equivalent years of full health lost through living in poor health from birth is also greater in women than in men (9.5 versus 7.8 years).

Fig. O.1
Global life expectancy and healthy life expectancy, 2000–2016




Source: Global health estimates 2016: Life expectancy, 2000–2016. Geneva: World Health Organization; 2018 (https://www.who.int/gho/mortality_burden_disease/life_tables/en/, accessed 21 March 2019).

The ratio of the number of males alive to the number of females alive changes through the life-course

Globally, the sex ratio at birth has been in the range of 105–110 males to every 100 females; however, mortality rates are higher in males, so the ratio changes as the population ages. Thus, in 2016, there were 100 men for every 100 women in the age group 50–54 years, and 95 for every 100 women in the age group 60–64 years, with the sex ratio falling sharply thereafter (Fig. 0.2). Because the incidence of different diseases varies with age, and women live longer than men, some diseases can be more common in women; for example, the lifetime risk for Alzheimer’s disease is greater in women than in men, partly because more women survive to ages at which the disease most commonly occurs, although in some locations women also appear to be more susceptible to Alzheimer’s disease (2).

Fig. 0.2
Global male to female (M/F) sex ratio by age, 2016



Source: World population prospects 2017 (3).


Several conditions contribute to differences in life expectancy between men and women

The reduced life expectancy of males compared with that of females is not due to a single or a small number of causes. Of the 40 leading causes of death, 33 contribute more to reduced life expectancy in males than in females (Fig. 0.3). The main causes of death that contribute to a lower life expectancy in males than in females are ischaemic heart disease (0.84 years), road injuries (0.47), lung cancers (0.40), chronic obstructive pulmonary disease (0.36), stroke (0.32), cirrhosis of the liver (0.27), tuberculosis (TB) (0.23), prostate cancer (0.22) and interpersonal violence (0.21).

Breast cancer (0.30 years), maternal conditions (0.23) and cervical cancer (0.15) are the causes of death that most reduce female rather than male global life expectancy.

Some of the differences in mortality rates and life expectancy are due to biological sex differences between females and males. For example, X-linked immune regulators may enhance immune responses in female children, resulting in reduced mortality among girls aged under 5 years. Others are linked to gender differences; that is, the socially constructed roles, norms, behaviours, activities and attributes that a given society considers appropriate for men, women boys and girls. For example, child marriage increases the risks related to early pregnancy among girls, whereas higher rates of male employment in the transport industry expose men to higher risks of death on the roads. The exact contributions that biological differences and gender roles make to health status are often difficult to determine because they do not operate independently (4).

Figure 0.3
Causes of death that most contribute to differences in life expectancy at birth globally for men and women, 2016



Source: WHO global health estimates 2016 (1), and see (5) for decomposition of life expectancy.

Life expectancy and age of death varies greatly by country income group

Life expectancy at birth in low-income countries (62.7 years) is 18.1 years lower than in high-income countries (80.8 years) (Table O.2). In high-income countries, most of the people who die are old; however, in low-income countries almost one in three deaths are of children aged under 5 years (Fig. O.4).

Table O.2
Life expectancy and HALE by sex, WHO region and World Bank income group, 2016

		Life expectancy	HALE	Life expectancy	HALE
		At birth (years)		At 60 years (years)	
		Global	Male	69.8	62.0
	Female	74.2	64.8	21.9	16.8
	Both sexes	72.0	63.3	20.5	15.8
WHO region (both sexes)	AFR	61.2	53.8	16.6	12.5
	AMR	76.8	67.5	22.7	17.6
	SEAR	69.5	60.4	18.2	13.3
	EUR	77.5	68.4	22.3	17.4
	EMR	69.1	59.7	18.2	13.3
	WPR	76.9	68.9	21.0	16.6
World Bank income group (both sexes)	LI	62.7	54.9	17.1	12.9
	LMI	67.9	59.1	18.0	13.2
	UMI	75.2	67.0	20.2	15.8
	HI	80.8	71.2	24.3	19.0

Darker shading represents lower values of life expectancy.


Source: Global health estimates 2016: Life expectancy, 2000–2016. Geneva: World Health Organization; 2018 (https://www.who.int/gho/mortality_burden_disease/life_tables/en/, accessed 21 March 2019). World Bank classification of economies July 2017.

Differences in life expectancy between men and women are greater in higher income countries

The differences in life expectancy between females and males are smaller in low-income countries than in high-income countries. Communicable diseases, injuries and maternal conditions contribute most to differences in life expectancy between females and males in low-income countries, whereas noncommunicable diseases (NCDs) contribute most to life expectancy differences in high-income countries (Fig. O.5).


Maternal deaths contribute more than any other cause to differences in life expectancy at birth between men and women. Maternal deaths are concentrated in low-income countries, being related primarily to lack of access to essential health services. The life expectancy of men is lower than that of women due to higher mortality rates from most causes, particularly in higher income countries; in low-income countries, the net effect of maternal conditions, breast and cervical cancer reduces the differences in life expectancy between men and women compared with high-income countries.

Fig. O.4
Proportion of deaths by age and World Bank income group, 2016 (%)



Source: World population prospects 2017 (3). World Bank classification of economies July 2017.

Fig. 0.5
Causes of death responsible for differences in life expectancy between men and women and World Bank income group, 2016



Source: WHO global health estimates 2016 (1). World Bank classification of economies July 2017.


The responses to differences in health status will vary according to the circumstances of countries

In low-income countries, differences in life expectancy between males and females are due to causes that are frequently preventable, or treatable through access to basic health services. In higher income countries, premature deaths are frequently associated with environmental factors or unhealthy lifestyle (Fig. 0.6).

Where there are differences in health outcomes between men, women, boys, girls and transgender people, gender-sensitive responses that aim to influence differences in exposure to risk factors or to increase access to health services may help to both minimize differences in health outcomes and maximize health gains.

Many of the factors that influence exposure to risks and access to health services are beyond the traditional remit of ministries of health. They require a multisectoral approach that addresses the underlying causes of gender and socioeconomic inequalities.

Fig. 0.6
Concentration of deaths according to national income of countries and sex, 2016



Points represent the 40 leading causes of death globally, with their areas being proportional to the number of deaths in 2016. Selected causes are labelled; space does not permit labelling of all causes. The concentration index is used to summarize the extent to which deaths from a disease are concentrated in high- or low-income countries, or in males or females. The index ranges from -1 to 1: a value of 0 indicates no association with national income or sex, and a value of -1 or 1 indicates that a disease occurs exclusively in males or females or in low-income or high-income countries (e.g. maternal deaths occur exclusively in women and are concentrated in low-income countries, whereas lung cancer is concentrated in high-income countries and occurs more in males).

Source: WHO global health estimates 2016 (1), and see (6) for calculation of concentration index. World Bank classification of economies July 2017.

HEALTH-RELATED SDG INDICATORS

3.1. Global trends

Globally, there have been improvements in most of the health-related SDG indicators

Recent years have seen improvements in 24 (56%) of the 43 health-related SDG indicators tracked in *World health statistics 2019*¹ (Table O.3). However, at a global level, progress has stalled or trends are in the wrong direction, for five of those 43 indicators: road traffic mortality, children overweight, malaria incidence, alcohol consumption, water

Table O.3
Trends in health-related SDG indicators

Programme area

- Reproductive, maternal and child health
- Infectious and noncommunicable diseases
- Injuries, violence and environmental risks
- Health systems and financing

SDG indicators with explicit targets for 2030

Progress stalled or trend in wrong direction	
3.6.1	Road traffic mortality
6.a.1	Water sector ODA
Progress made but too slow to meet target	
3.1.1	Maternal mortality
3.4.1	NCD mortality
6.1.1	Safe drinking-water coverage
6.2.1	Safe sanitation coverage
7.1.2	Clean energy coverage
Progress fast enough to attain target	
3.2.1	Under-5 mortality
3.2.2	Neonatal mortality

¹ Four of the health-related SDG indicators in the official list of SDG indicators (7) have more than one component (Indicators 2.2.2, 3.8.2, 3.b.1, 3.c.1). Where this is the case, each component is treated as a separate indicator in *World health statistics 2019*. The 43 indicators tracked have 36 unique indicator numbers in the official list of SDG indicators.

SDG indicators with no explicit targets for 2030

Progress stalled or trend in wrong direction	
2.2.2	Children overweight
3.3.3	Malaria incidence
3.5.2	Alcohol consumption
Progress made	
3.1.2	Skilled birth attendance
3.7.1	Met need for family planning
3.7.2	Adolescent birth rate
2.2.1	Stunting in children
3.b.1	DTP3 coverage
	MCV2 coverage
	PCV3 coverage
3.3.1	New HIV infections
3.3.2	Tuberculosis incidence
3.3.4	Hepatitis B prevalence
3.3.5	Need for NTD interventions
3.4.2	Suicide mortality
3.a.1	Tobacco use in persons ≥15 years
16.1.1	Homicide
3.9.3	Poisoning mortality
3.b.2	ODA med research & basic health
1.a.2	Government health expenditure

Trend not yet reported	
2.2.2	Wasting in children
3.9.1	Air pollution mortality
3.9.2	Unsafe water and sanitation mortality
5.2.1	Intimate partner violence
11.6.2	Fine particulate matter in urban areas
3.8.1	UHC service coverage index
3.c.1	Medical doctor density
	Nurse/midwife density
	Dentist density
	Pharmacist density
3.d.1	International Health Regulations capacity
3.8.2	Household health expenditures >10%
	Household health expenditures >25%
17.9.2	Completeness of cause-of-death data

sector ODA, and catastrophic health spending. Trends have not yet been reported in 14 of the 43 indicators (33%). Eight of the health-related SDG indicators have explicit targets for 2030, but only two of those indicators are on track to meet 2030 targets; that is, those for under-5 mortality rate and neonatal mortality rate. Moreover, it is estimated that on current trends 51 countries will miss the target for under-5 mortality, and more than 60 countries will miss the target for neonatal mortality in 2030.

3.2. Differences among countries

SDG indicators of health status

The disparities in life expectancy among countries are reflected in many of the health-related SDG indicators (Table O.4). Maternal mortality ratios are 29 times higher in low-income countries than in high-income countries. In resource-poor settings, fertility rates are higher and the risks

Table O.4
Latest values of health-related SDG indicators by sex, WHO region and World Bank income group

SDG indicator (by topic area)	Year	Global		WHO Region							WB Income group				2030 numeric target
		Male	Female	Both sexes	AFR	AMR	SEAR	EUR	EMR	WPR	LI	LMU	UMI	HI	
Reproductive and maternal health															
3.1.1 Maternal mortality ratio (per 1000 live births)	2015	-	216	-	542	52	164	16	166	41	495	253	55	17	<70
3.1.2 Proportion of births attended by skilled health personnel (%)	2013–2018	-	81	-	59	95	81	99	79	97	60	76	99	99	-
3.7.1 Family planning satisfied with modern methods ^c (%)	2019	-	76	-	56	83	71	77	61	87	-	-	-	-	
3.7.2 Adolescent birth rate (per 1000 women aged 15–19 years)	2015–2020	-	44	-	49	49	33	17	45	14	97	46	29	12	-
Child health															
3.2.1 Under-five mortality rate (per 1000 live births)	2017	41	37	39	74	14	36	9	50	13	69	49	14	5	25
3.2.2 Neonatal mortality rate (per 1000 live births)	2017	-	-	18	27	8	21	5	27	6	48	49	26	7	12
2.2.1 Prevalence of stunting in children under 5 (%)	2018	-	-	21.9	33.1	6.5	31.9	-	24.7	6.4	34.2	31.1	6.3	3.0	-
2.2.2 Prevalence of wasting in children under 5 (%)	2018	-	-	7.3	7.0	0.8	15.0	-	7.8	2.2	7.4	11.6	1.8	0.6	-
2.2.2 Prevalence of overweight in children under 5 (%)	2018	-	-	6.0	3.5	7.2	3.8	-	5.7	6.0	3.1	3.9	7.4	7.2	-
3.b.1 DTP3 immunization coverage among 1-year-olds (%)	2017	-	-	85	72	91	88	94	81	97	78	82	94	95	-
3.b.1 MCV3 immunization coverage by the nationally recommended age (%)	2017	-	-	67	25	74	77	90	67	94	29	63	88	91	-
3.b.1 PCV3 immunization coverage among 1-year olds (%)	2017	-	-	44	68	82	12	70	52	16	68	32	33	85	-
Infectious diseases															
3.3.1 New HIV infections (per 1000 uninfected population)	2017	0.26	0.24	0.25	1.22	0.16	0.08	0.18	0.06	0.06	0.66	0.23	0.24	0.07	-
3.3.2 Tuberculosis incidence (per 100 000 population)	2017	168	99	133	237	28	226	30	113	94	244	223	58	11	-
3.3.3 Malaria incidence (per 1000 population at risk)	2017	-	-	59.1	219.4	7.3	7.0	-	14.8	2.5	189.3	42.8	2.5	-	-
3.3.4 Hepatitis B surface antigen prevalence among children under 5 years (%)	2017	-	-	0.80	2.34	0.07	0.26	0.21	0.69	0.38	2.31	0.72	0.30	0.16	-
3.3.5 Reported number of people requiring interventions against NTDs (millions)	2017	-	-	1583	594	76	733	5	75	98	398	1069	115	<1	-
Noncommunicable diseases															
3.4.1 Probability of dying from CVD, cancer, diabetes, CRO between age 30 and exact age 70 (%)	2016	21.6	15.0	18.3	20.6	15.1	23.1	16.7	22	16.2	21.3	23.3	17.7	12.0	Reduce 1/3
3.4.2 Suicide mortality rate (per 100 000 population)	2016	13.5	7.7	10.6	7.4	9.8	13.2	15.4	3.9	10.2	6.8	10.6	10.0	14.3	Reduce 1/3
3.5.2 Total alcohol per capita (≥15 years of age) consumption (limes of pure alcohol)	2016	10.1	2.7	6.4	6.3	8.0	4.5	9.8	0.6	7.3	-	-	-	-	-
3.a.1 Prevalence of tobacco smoking among persons aged 15 years and older ^d (%)	2016	33.7	6.2	19.9	9.8	16.9	16.9	29.4	18.1	24.5	11.4	17.2	23.1	24.1	-
Injuries and violence															
3.6.1 Road traffic mortality rate (per 100 000 population)	2016	-	-	18.2	26.6	15.6	20.7	9.3	18	16.9	27.5	19.2	8.3	8.3	Half by 2020
5.2.1 Proportion of women aged 15 years and older subjected to intimate partner violence (%)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
16.1.1 Mortality rate due to homicide (per 100 000 population)	2016	10.1	2.6	6.4	10.4	17.9	4.1	3.3	6.7	1.9	8.7	5.9	7.9	2.9	-
Environmental risks															
3.9.1 Mortality rate attributed to household and ambient air pollution ^b (per 100 000 population)	2016	128.5	101.1	114.1	180.9	29.7	165.8	36.3	125	102.8	-	131.7	-	17.8	-
3.9.2 Mortality rate attributed to exposure to unsafe WASH services (per 100 000 population)	2016	11.4	12.1	11.7	45.8	1.1	15.4	0.3	10.6	1.0	42.4	18.6	1.1	0.3	-
3.9.3 Mortality rate from unintentional poisoning (per 100 000 population)	2016	1.6	1.2	1.4	2.7	0.6	1.8	0.7	1.5	1.1	2.8	1.8	1.1	0.5	-
6.1.1 Proportion of population using safely managed drinking-water services (%)	2015	-	-	71	26	82	-	91	56	-	23	59	92	98	Universal
6.2.1 Proportion of population using safely managed sanitation services (%)	2015	-	-	39	-	43	-	67	-	57	-	-	50	81	Universal
6.a.1 WASH-related ODA ^e (constant 2016 US\$ millions)	2017	-	-	8698.25	2483.89	676.69	1484.41	635.52	1836.26	1011.10	1983.59	4262.35	1750.49	-	-
7.1.2 Proportion of population with primary reliance on clean fuels (%)	2017	-	-	60	17	92	42	97	72	64	52	99	99	99	Universal
11.6.2 Annual mean concentrations of fine particulate matter (PM2.5) in urban areas (µg/m3)	2016	-	-	39.6	35.5	13.4	57.3	17.6	54	42.9	44.0	-	-	14.4	-
UHC and health systems															
3.8.1 UHC service coverage index	2015	-	-	64	44	78	55	73	53	75	40	54	74	80	Universal
3.8.2 Catastrophic out-of-pocket health spending > 10% ^a	2010	-	-	11.7	10.3	11.1	12.8	7.0	9.5	14.8	8.1	12.4	13.8	7.2	Universal
3.8.2 Catastrophic out-of-pocket health spending > 25% ^a	2010	-	-	2.6	2.6	1.9	2.8	1.0	1.4	3.9	1.1	2.8	3.2	1.4	Universal
1.a.2 Domestic general government health expenditure as percentage of GGE (%)	2016	-	-	10.6	7.3	15.6	6.7	12.5	8.5	11	6.6	8.1	11.5	14.9	-
3.c.1 Density of medical doctors (per 10 000 population)	2017	-	-	15.1	2.8	23.3	7.4	33.8	9.9	18	3.1	7.5	19.4	30.4	-
3.c.1 Density of nursing and midwifery personnel (per 10 000 population)	2017	-	-	34.8	10.9	61.9	19.9	80.6	15.2	32.6	8.5	18.9	35	85.6	-
3.c.1 Density of dentists (per 10 000 population)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
3.c.1 Density of pharmacists (per 10 000 population)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
3.d.1 Average of 13 International Health Regulations core capacity scores	2018	-	-	60	42	64	56	72	66	62	42	52	63	76	-
3.b.2 Total net ODA to medical research and basic health sectors per capita (US\$)	2017	-	-	1.39	4.83	0.42	0.60	-	1.89	0.30	5.64	1.23	0.33	-	-
17.19.2 Completeness of cause-of-death data (%)	2017	-	-	49	6	94	10	97	32	64	3	15	73	97	-


- Not available or applicable
- a Women of reproductive age
- b Age-standardized
- c Ever-partnered women and girls aged 15 years and older subjected to physical, sexual or psychological violence by a current or former intimate partner in the previous 12 months
- d Amount that is part of a government-coordinated spending plan, refers to water sector only
- e Population with household expenditures on health greater than 10 or 25% of total household expenditure or income
- f recipient countries
- GGE general government expenditure

Darker shading represents high values of indicators for mortality, incidence, prevalence and risk indicators; and lower values of indicators for programme coverage and financing indicators.

Source: World Health Statistics 2019 (forthcoming).

of dying in labour greater, so the lifetime risk of maternal death is greatly amplified; in low-income countries, one woman out of 41 dies from maternal causes (Fig. O.7).

Fig. O.7
Lifetime risk of dying from maternal causes by World Bank income group and WHO region, 2015



The life time risk of dying from maternal causes is the probability of a 15-year-old girl eventually dying from a maternal cause, assuming that she is subjected throughout her lifetime to the fertility and maternal mortality risks, as estimated for 2015.

Source: *Trends in maternal mortality: 1990 to 2015 (8)*.


In low-income countries, about a third of children are stunted (i.e. short for their age), reflecting long-term nutritional deprivation, and one child out of 14 will die before their fifth birthday. Adolescent birth rates are eight times higher in low-income countries than in high-income countries. The burden of infectious diseases, including HIV, TB, malaria, hepatitis B and neglected tropical diseases (NTDs) is higher in low-income countries than in high-income countries. Mortality rates attributed to unsafe water, unsafe sanitation and lack of hygiene are also highest in low-income countries, as are mortality rates for road traffic injuries and unintentional poisoning. Although NCDs are often associated with a more prosperous lifestyle, the probability of dying prematurely from cardiovascular disease (CVD), cancer, diabetes and chronic respiratory disease is highest in low- and lower-middle-income countries.

There are some exceptions to poor health being primarily associated with low-income countries. Highest wasting rates are observed in the WHO South-East Asia Region (15.0%) and Eastern Mediterranean Region (7.8%). Mortality rates from homicide are highest in the WHO Region of the Americas. Suicide mortality rates are highest in the WHO European Region, and lowest in the Eastern Mediterranean Region.

SDG indicators of health service coverage and financing

Populations in low-income countries generally have less access to essential health services; values of the universal health coverage (UHC) service coverage index are lower, as are indicators such as skilled birth attendance, women who have their need for family planning satisfied with modern methods of contraception, and immunization coverage. Low-income countries also experience greater shortages of health care professionals (Fig. O.8), and domestic government health expenditure as a proportion of total general government expenditures is lower (despite lower absolute levels of general government expenditure and greater health needs). At all income levels people can suffer catastrophic health expenditures (out-of-pocket health spending >10% or 25% of total household expenditures or income), even in high income countries and in countries where most of the out-of-pocket health spending is due to medicines.

Fig. O.8
Percentage of countries with insufficient health care professionals, 2013–2018



Source: WHO Global Health Workforce Statistics [online database], Global Health Observatory (GHO) data. Geneva: World Health Organization (<https://who.int/hrh/statistics/hwfstats/en/>, accessed 15 March 2019).

SDG indicators of exposure to risk factors.


Populations in lower-income countries are less likely to use safely managed drinking-water, and clean fuels and technology; also, they have greater exposure to fine particulate matter in cities. In contrast, tobacco use and alcohol consumption are highest in high-income countries.

3.3. Differences between females and males

SDG indicators of health status

Differences between females and males are seen in most of the health-related SDG indicators for which sex disaggregation has been possible. In 2017, male children were 11% more likely to die than female children before the age of 5 years, compared with only 6% in 2000, indicating that the decline in under-5 mortality rate since 2000 has been faster in females than in males. Given that boys generally have a higher under-5 mortality rate than girls for biological reasons, the similarity of the under-5 mortality rate between boys and girls in the WHO South-East Asia Region is indicative of high rates of avoidable mortality among female children. Although global estimates are not available, household surveys indicate

Fig. 0.9
Mainnutrition rates in female and male children: (a) stunting (low height for age), (b) wasting (low weight for height)




Each point represents latest survey values available for a county 2000–2017.
 Source: Levels and trends in child malnutrition. UNICEF/WHO/World Bank Joint Child Malnutrition Estimates 2019 Edition, available at <http://www.who.int/nutgrowthdb/estimates>. World Bank classification of economies July 2018.

that rates of stunting and wasting are generally higher in boys than in girls (Fig. O.9). Similarly, the proportion of boys overweight is higher than the proportion of girls.

Globally in 2017, the incidence rate of new HIV infections was 1.09 times as high as in men than in women. In sub-Saharan Africa, the incidence was 1.27 times as high as in women than in men; however, in other parts of the world, the incidence was 1.7 times as high as in men than women, reflecting differences in predominant modes of transmission. The incidence rate of TB rates were 1.7 higher in men than in women globally in 2017. In 2016, the probability of a man aged 30 years dying from an NCD before 70 years of age was 1.44 times higher than for a woman aged 30 years (Fig. O.10). Globally in 2016, suicide mortality rates were 1.75 times higher in men than in women.


Fig. O.10
Age-standardized rates of premature death from the four major NCDs covered by SDG Target 3.4, 2016



Source: WHO global health estimates 2016 (1).

Deaths rates from road injury are more than twice as high in men as in women from age 15 years (Fig. O.11), and mortality rates due to homicide are four times higher in men than in women. One in five of all homicides is committed by an intimate partner or family member, with women making up most of the victims (9). Beyond mortality, violence against women is not only widespread, but carries a high burden of morbidity and ill health. Worldwide, in 2013, it was estimated that 35% of women and girls aged 15–49 years reported physical or sexual intimate partner violence or non-partner sexual violence in their lifetime. Women and girls may also experience harmful practices, such as female genital mutilation (FGM) and early and forced marriage. As of 2017, at least 200 million girls and women had undergone FGM in the 30 countries where the practice is concentrated (10).

Fig. O.11
Ratio of global M/F crude death rates by age and cause, 2016



Source: WHO global health estimates 2016 (1). Death rates for road traffic injury from this source may differ from figures presented elsewhere in this report, but are used in this graph for comparison with other causes of injury.

Globally, 21% of women aged between 20 and 24 years reported that they were married or in an informal union before age 18 years (11). Also, in 2015–2020, it is estimated that more than one in 25 adolescent girls aged 15–19 years will give birth (3).


Mortality rates attributed to household and ambient air pollution and unintentional poisoning are 1.27 times higher in men than in women. In contrast, the global mortality rate

attributed to unsafe water, unsafe sanitation and lack of hygiene is 1.06 higher in women than in men, although there is much regional variation.

SDG indicators of health service coverage and financing

Access to many services for prevention or treatment of infectious diseases and NCDs is not part of the SDG monitoring framework, except as part of the UHC service coverage index, which cannot be sex-disaggregated using currently available data and methodologies. Nevertheless, some relevant observations have been made through other global health reports. Household surveys suggest that vaccination rates are similar in boys and girls. The risk of not using a condom during sex with a non-regular partner appears to be higher in women than in men. However, in countries with generalized HIV epidemics, men are less likely than women to take an HIV test and less likely to access antiretroviral therapy; also, men are more likely than women to die of an AIDS-related illness (12). Similarly, male TB patients appear to be less likely to seek care than female TB patients (13–16), as reflected in lower rates of case notification compared with the estimated total of cases (Fig. O.12). Death rates for some NCDs may also be influenced by access to diagnosis and treatment; for example, cervical cancer mortality rates are higher in low-income countries that have

Fig. O.12
Estimated incidence of TB in comparison to case notification by World Bank income group, age and sex, 2017



Source: Global tuberculosis report 2018 (16). World Bank classification of economies July 2018


poorer access to health services.

SDG indicators of exposure to risk factors

Age-standardized prevalence of tobacco smoking was five times higher in men than in women in 2016, with the largest M/F ratio observed in the WHO Eastern Mediterranean Region (Fig. O.13). Globally, per capita alcohol consumption was almost 4 times higher in men than in women in 2016.

Fig. O.13

Among persons aged 15 years and older: (a) age-standardized prevalence of tobacco smoking, 2016, (b) alcohol consumption per capita, 2016



Source: World Bank classification of economies July 2017; a) WHO global report on trends in prevalence of tobacco smoking, 2nd edition. Geneva: World Health Organization; 2018 (<https://www.who.int/tobacco/publications/surveillance/trends-tobacco-smoking-second-edition/en/>, accessed 15 March 2019).

Source: World Bank classification of economies July 2017; b) WHO Global Information System on Alcohol and Health (GISAH) [online database], Global Health Observatory (GHO) data. Geneva: World Health Organization (<https://www.who.int/gho/alcohol/en/>, accessed 15 March 2019).

4 DATA FOR TRACKING THE HEALTH-RELATED SDGs


Monitoring of the health-related SDGs is based on statistics of two types:

- **primary data** – data compiled by international agencies from routine reporting by countries or publicly available sources such as demographic and health surveys; statistics are presented as they are reported and no (or minimal) adjustments are made to values; and
- **comparable estimates** – country data are adjusted or modelled to allow comparisons among countries or over time.

For 18 SDG indicators that are reported as primary data, the proportion of countries with available primary data within the past 10 years ranged from 35% for intimate partner violence, to 100% for International Health Regulations (2005) core capacity scores and need for NTD interventions (Fig. O.14). The proportion of countries with available primary data within the past 5 years ranged from 5% to 100%.

For 25 SDG indicators that are reported as comparable estimates, the availability of underlying data also varies across indicators (Fig. O.15). Indicators that were adopted for global monitoring before the SDG era – for example,

Fig. O.14
The availability of underlying data for SDG indicators reported as primary data



a) Denominator for 3.b.2 and 6.a.1 reflects the list of official development assistance (ODA) recipients 2014–2017 maintained by the Development Assistance Committee of the Organisation for Economic Co-operation and Development.

b) Indicator 3.8.2 considers two thresholds (10% and 25% of total household expenditure); availability the same for both. Indicator 3.c.1, health workforce, includes four components (density of medical doctors, nursing and midwifery personnel, dentists and pharmacists); availability is shown for density of nursing and midwifery personnel. Indicator 2.2.2 includes two components (proportion of children wasted and proportion of children overweight); availability shown for children wasted.

Fig. 0.15
The availability of underlying data for SDG 3 indicators reported as comparable estimates



a) Denominator for Indicator 3.3.3 is the number of endemic countries in 2000 (107).

b) Mortality attributed to household and ambient air pollution, unsafe WASH services and unintentional poisoning.

under-5 mortality rate, TB and HIV incidence, and vaccination coverage – tend to have relatively high availability of underlying data, even though such data may not be recent for all countries. In contrast, availability of underlying data is lower for new global indicators such as cause-specific mortality rates, and proportion of population using safely managed drinking-water and safely managed sanitation services. The average lag between the year of the estimate and the latest available year of underlying data ranged from 0 years for HIV, TB and malaria incidence, immunizations, road traffic mortality and UHC service coverage index to 5 years for hepatitis B prevalence.

Of the 43 health and health-related SDG indicators reported in *World health statistics 2019*, sex disaggregation would potentially be informative for 28 indicators (Table O.5). Sex disaggregation is not possible for the ten indicators for which data are collected at household or national level, or for five indicators that have female-specific numerators or denominators. For the 28 SDG indicators for which sex-disaggregated indicators could be informative, sex-disaggregated global and regional values are currently available for only 11 indicators (though sex-disaggregated values may be available at country level).

Table O.5
Availability of sex-disaggregated values of SDG indicators at global or regional levels

Programme area: ■ Reproductive, maternal and child health ■ Infectious and noncommunicable diseases ■ Injuries, violence and environmental risks ■ Health systems and financing

Sex disaggregated values reported	Sex disaggregated values not currently reported	Data collected at household or higher level
3.2.1 Under-5 mortality	3.2.2 Neonatal mortality	6.1.1 Safe drinking-water coverage
3.3.1 New HIV infections	2.2.1 Stunting in children	6.2.1 Safe sanitation coverage
3.3.2 Tuberculosis incidence	2.2.2 Wasting in children	6.a.1 Water sector ODA
3.4.1 NCD mortality	2.2.2 Overweight children	7.1.2 Clean energy coverage
3.a.1 Tobacco use in persons ≥15 years	3.b.1 DTP3 immunization coverage	11.6.2 Fine particulate matter in urban areas
3.5.2 Alcohol consumption	MCV2 immunization coverage	3.8.2 Household health expenditure >10%
3.4.2 Suicide mortality	PCV3 immunization coverage	Household health expenditure >25%
16.1.1 Homicide	3.3.3 Malaria incidence	1.a.2 Government health expenditure
3.9.3 Poisoning mortality	3.3.4 Hepatitis B prevalence	3.b.2 ODA med research & basic health
3.9.1 Air pollution mortality	3.3.5 Need for NTD interventions	3.d.1 International Health Regulations capacity
3.9.2 Unsafe water and sanitation mortality	3.6.1 Road traffic mortality	
	3.8.1 UHC service coverage index	
	3.c.1 Medical doctor density	
	Nurse/midwife density	
	Dentist density	
	Pharmacist density	
	17.9.2 Cause-of-death data completeness	
		Female specific
		3.7.1 Met need for family planning
		3.7.2 Adolescent birth rate
		3.1.1 Maternal mortality
		3.1.2 Skilled attendance at birth
		5.1.2 Intimate partner violence



CONCLUSIONS

Some findings are evident from this report, even though more work is needed for a better understanding of the factors influencing risk behaviours, access to services and health outcomes. The largest disparities in life expectancy – and values of SDG indicators – occur when comparing country groupings such as WHO regions or World Bank income groups. Differences in the values of life expectancy and SDG indicators between men and women are generally smaller. Even so, there are some key areas where differences between men and women are pronounced, and where failure to address these differences will not only allow inequities to persist but could also jeopardize the attainment of health-related SDGs. Those areas are as follows:

- Under-5 mortality rates are generally higher in boys than girls owing to a greater biological frailty in boys. Under-5 mortality rates that are similar or equal between boys and girls are indicative of avoidable mortality among female children. As under-5 mortality rates fall globally, particularly in girls, the M/F mortality ratio is increasing. Hence, it may be necessary to consider additional actions that could be taken to improve health outcomes for boys, to ensure continued progress towards SDG Target 3.2.
- In low-income countries, maternal deaths contribute more to differences in life expectancy between men and women than any other single cause, whereas such deaths rarely occur in upper-middle and high-income countries. Maternal deaths can be reduced by ensuring that women have access to high-quality care before, during and after childbirth, and access to modern methods of contraception. Cervical cancer is also concentrated in low-income and lower-middle-income countries, and mortality rates can be substantially reduced through better access to health services including vaccination, screening and treatment.
- Men's reduced life expectancy compared with that of women is due to higher death rates from multiple causes, especially CVD, road injuries, lung cancers, chronic obstructive pulmonary disease and stroke. Better access to health services, such as treatment for high blood pressure, can increase survival rates for some of these conditions, but great impact can be made by altering exposure to risk factors such as tobacco, alcohol, unhealthy diet and road accidents. Although these risk factors are also important for women, both men and women could benefit from more gender-sensitive targeting of behavioural and legislative interventions to reduce exposure to harmful practices.
- In many settings, men use health services less than women, even after taking into account reproductive-related consultations. For some infectious diseases, such as HIV and TB, men's later diagnosis and treatment jeopardizes the men's health and contributes to a greater number of secondary infections. Therefore, strategies to improve men's access to and use of health services for infectious diseases can not only help to address gender disparities in health outcomes, it can also help to reduce disease incidence in the general population. Although women may use service more than men it cannot be concluded that their use is optimal. Women often have less control over decision-making. Men may dominate decisions about sexuality and pregnancy, and access to sexual and reproductive health information or services; also, they may have greater control than women over household resources.
- Women are the major contributors to delivery of health services, but women are more likely to be in caring roles (often unpaid) and less likely to be employed as, for example, doctors or managers. The under-representation of women as managers may lead to a poor understanding of working conditions of the predominantly female workforce, and of health care needs specific to women.



WAYS FORWARD

Some key actions are suggested based on the findings of this report:

1. Develop policies and programmes that are sensitive to sex and gender

Health planning should take into account differences between females and males in exposure to risk factors, access to services and differences in health outcomes, and should be prepared to target actions to areas where sex or gender differences contribute most to health outcomes. Gender analysis should consider women, men and gender-diverse population groups to ensure equitable health outcomes.

2. Invest in data systems for health, including sex-disaggregated data

The development of policies and programmes to address inequities in health requires better information on differences between men and women, and other dimensions of inequity such as socioeconomic status and place of residence. This may require modifications to the way routine information systems and health facility or household surveys are implemented, and the way that data are reported and analysed. It will also require strengthening of civil vital registration systems. Qualitative research is also needed that can reveal social and cultural factors underlying gender norms, roles, relations and expectations that lead to poor health outcomes.

3. Build responsive institutions

To enable decent conditions of work for all and responsive health systems, it is necessary to formulate gender-sensitive human resource policies, ensure equal pay for work of equal value, and address barriers faced by women such as mobility restrictions and lack of education opportunities. There is also a need for increased leadership roles for women.

4. Address upstream influences on health status and use of health services

Many of the factors that influence health and use of health services are beyond the traditional remit of ministries of health. Multisectoral approaches are needed that address underlying factors, such as gender and socioeconomic inequalities. Nonetheless, the health sector has a key role in raising awareness of these issues, and can catalyse the development of gender-sensitive and gender-transformative policies and programmes.

5. Consider gender differences that are not limited to female disadvantage

In many circumstances, men experience poorer health outcomes than women do. Although some of these poorer health outcomes may have a biological basis, they may be amplified by gender roles. Health policies and programmes need to be responsive to the needs of both men and women.

REFERENCES

1. Global health estimates 2016: deaths by cause, age, sex, by country and by region, 2000–2016. Geneva: World Health Organization; 2018.
2. Mielke MM, Ferretti MT, Iulita MF, Hayden K, Khachaturian AS. Sex and gender in Alzheimer's disease – does it matter? *Alzheimer's & Dementia*. 2018;14(9):1101 (<https://www.ncbi.nlm.nih.gov/pubmed/30196887>, accessed 26 March 2019).
3. World population prospects. The 2017 revision. Geneva: United Nations, Department of Economic and Social Affairs, Population Division; 2017 (<https://population.un.org/wpp/>, accessed 31 March 2019).
4. Hawkes S, Buse K. Gender and global health: evidence, policy, and inconvenient truths. *Lancet*. 2013;381(9879):1783–7.
5. Beltran-Sanchez H, Preston SH, Canudas-Romo V. An integrated approach to cause-of-death analysis: cause-deleted life tables and decompositions of life expectancy. *Demogr Res*. 2008;19:1323 (<https://www.ncbi.nlm.nih.gov/pubmed/20165568>, accessed 31 April 2019).
6. O'Donnell O, van Doorslaer E, Wagstaff A, Lindelow M, Analyzing Health Equity Using Household Survey Data. The World Bank, Washington DC, 2008, .
7. SDG Indicators: Global indicator framework for the Sustainable Development Goals and targets of the 2030 Agenda for Sustainable Development. [website]. United Nations; (<https://unstats.un.org/sdgs/indicators/indicators-list/>, accessed 31 April 2019).
8. Trends in maternal mortality: 1990 to 2015: estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division. Geneva: World Health Organization; 2015 (https://apps.who.int/iris/bitstream/handle/10665/194254/9789241565141_eng.pdf;jsessionid=4F32A9EABB0E2D5B373778396FEB6EB?sequence=1, accessed 31 April 2019).
9. https://www.unodc.org/documents/data-and-analysis/GSH2018/GSH18_Gender-related_killing_of_women_and_girls.pdf
10. <https://data.unicef.org/topic/child-protection/female-genital-mutilation/>
11. <https://data.unicef.org/wp-content/uploads/2018/07/Child-Marriage-Data-Brief.pdf>
12. Blind spot: addressing a blind spot in the response to HIV. Geneva: Joint United Nations Programme on HIV/AIDS; 2017 (http://www.unaids.org/en/resources/documents/2017/blind_spot, accessed 17 March 2019).
13. Amere GA, Nayak P, Salindri AD, Narayan KMV, Magee MJ. Contribution of smoking to tuberculosis incidence and mortality in high-tuberculosis-burden countries. *Am J Epidemiol*. 2018;187(9):1846–55 (<http://dx.doi.org/10.1093/aje/kwy081>, accessed 17 March 2019).
14. Narasimhan P, Wood J, MacIntyre CR, Mathai D. Risk factors for tuberculosis. *J Pulm Med*. 2013;2013:1–11 (<http://dx.doi.org/10.1155/2013/828939>, accessed 17 March 2019).
15. Horton KC, MacPherson P, Houben RMGJ, White RG, Corbett EL. Sex differences in tuberculosis burden and notifications in low- and middle-income countries: a systematic review and meta-analysis. *PLOS Med*. 2016;13(9):e1002119
16. Global tuberculosis report 2018. Geneva: World Health Organization; 2018 (https://www.who.int/tb/publications/global_report/en/, accessed 31 August 2019).

Notes

A series of horizontal dotted lines for writing notes.

Notes

A series of horizontal dotted lines for writing notes.

Notes

A series of horizontal dotted lines for writing notes.

